

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 May 2004 (21.05.2004) ✓

PCT

(10) International Publication Number
WO 2004/041826 A1 ✓

(51) International Patent Classification⁷: **C07D 487/04**,
A61K 31/4985, 31/53, A61P 25/00 // (C07D 487/04,
241:00, 235:00) (C07D 487/04, 253:00, 235:00)

[GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex
CM20 2QR (GB). **STREET, Leslie, Joseph** [GB/GB];
Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR
(GB).

(21) International Application Number:

PCT/GB2003/004685 ✓

(74) **Agent: HORGAN, James, Michael, Fred**; European
Patent Department, Terlings Park, Eastwick Road, Harlow,
Essex CM20 2QR (GB).

(22) International Filing Date: 29 October 2003 (29.10.2003) ✓

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0225923.2	6 November 2002 (06.11.2002)	GB
0302529.3	4 February 2003 (04.02.2003)	GB
0304415.3	26 February 2003 (26.02.2003)	GB
0313646.2	12 June 2003 (12.06.2003)	GB

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) **Applicant (for all designated States except US): MERCK
SHARP & DOHME LIMITED** [GB/GB]; Hertford Road,
Hoddesdon, Hertfordshire EN11 9BU (GB).

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only): CARLING,
William, Robert** [GB/GB]; Terlings Park, Eastwick Road,
Harlow, Essex CM20 2QR (GB). **CASTRO PINEIRO,
Jose, Luis** [ES/GB]; Terlings Park, Eastwick Road,
Harlow, Essex CM20 2QR (GB). **GOODACRE, Simon,
Charles** [GB/GB]; Terlings Park, Eastwick Road, Harlow,
Essex CM20 2QR (GB). **HALLETT, David, James**


(84) **Designated States (regional):** ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) **Title: MIDAZOPYRAZINONES AND IMIDAZOTRIAZINONES DERIVATES AS GABA-A RECEPTOR ANXIOLYTIC**



(I)

(57) **Abstract:** The present invention discloses a compound of formula I, or a pharmaceutically acceptable salt thereof: (I) wherein -U-V- represents -CH=CH-, or -CH₂-CH₂-, -N=CH- or -CH=N-; X¹ represents hydrogen, halogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy; X² represents hydrogen or halogen; Y represents a chemical bond, an oxygen atom, or a -NH- or -OCH₂- linkage; Z represents an optionally substituted aryl or heteroaryl group; R₁ represents hydrocarbon, a heterocyclic group, trifluoromethyl, -SO₂R^a, -SO₂NR^aR^b, -COR^a, -CO₂R^a or -CONR^aR^b; and R^a and R^b independently represent hydrogen, hydrocarbon or a heterocyclic group; pharmaceutical compositions comprising it; its use in methods of treatment; use of it in the manufacture of a medicament for treating and/or preventing anxiety; convulsions or a cognitive disorder; and methods of treatment using it.

WO 2004/041826 A1

THIS PAGE BLANK (USPTO)


MIDAZOPYRAZINONE AND IMIDZOTRIAZINONE DERIVATIVES AS GABA-A RECEPTOR ANXIOLYTICS

The present invention relates to classes of substituted imidazo-pyrazinone derivatives, substituted imidazotriazine derivatives and purinone derivatives and to their use in therapy. More particularly, this invention is concerned with imidazo[1,2-*a*]pyrazin-8-one analogues and imidazo[1,2-*d*]triazin-8-one analogues which are substituted in the 3-position by a substituted phenyl ring, and with purin-6-one analogues which are substituted in the 9-position by a substituted phenyl ring. These compounds are ligands for GABA_A receptors and are therefore useful in the therapy of deleterious neurological complaints.

A description of the theoretical background underlying this invention can be found on pages 1-5 of WO-A-02074773 (Merck Sharp & Dohme Ltd.).

WO 02/10170 describes a class of 3-phenylimidazo[1,2-*a*]pyrazine derivatives which are stated to be selective ligands for GABA_A receptors, in particular having high affinity for the $\alpha 2$ and/or $\alpha 3$ subunit thereof, and accordingly to be of benefit in the treatment and/or prevention of neurological disorders, including anxiety and convulsions. However, there is no disclosure nor any suggestion in that publication of therapeutic agents based on a 7-substituted imidazo[1,2-*a*]pyrazin-8-one ring system.

The present invention provides a compound of formula I, or a pharmaceutically acceptable salt thereof:



(I)

25

wherein

-U-V- represents -CH=CH-, or -CH₂-CH₂-, -N=CH- or -CH=N-;

X¹ represents hydrogen, halogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy;

X² represents hydrogen or halogen;

Y represents a chemical bond, an oxygen atom, or a -NH- or -OCH₂- linkage;

5 Z represents an optionally substituted aryl or heteroaryl group;

R¹ represents hydrocarbon, a heterocyclic group, trifluoromethyl, -SO₂R^a, -SO₂NR^aR^b, -COR^a, -CO₂R^a or -CONR^aR^b; and

R^a and R^b independently represent hydrogen, hydrocarbon or a heterocyclic group.

10 The present invention also provides a compound of formula I as depicted above, or a pharmaceutically acceptable salt thereof, wherein

-U-V- represents -CH=CH-;

Y represents a chemical bond, an oxygen atom, or a -NH- linkage; and

X¹, X², Z and R¹ are as defined above.

15 Definitions of the aryl or heteroaryl group Z, pharmaceutically acceptable salts of the compounds, the terms "hydrocarbon" and "a heterocyclic group" as used herein, suitable alkyl, alkenyl, alkynyl, and cycloalkyl groups, typical examples of C₃₋₇ cycloalkyl(C₁₋₆)alkyl groups, particular indanyl, aryl and aryl(C₁₋₆)alkyl groups, suitable heterocycloalkyl and heteroaryl groups, the expression
20 "heteroaryl(C₁₋₆)alkyl", optional substituents on the hydrocarbon and heterocyclic groups and the term "halogen" as used herein can be found on pages 6-8 of WO-A-02074773.

General comments about stereochemistry can be found on page 9 of WO-A-02074773.

25 In one embodiment of the compounds according to the present invention, -U-V- represents -CH=CH-.

In another embodiment, -U-V- represents -CH₂-CH₂-.

Suitable values for the X¹ substituent include hydrogen, fluoro, chloro, methyl, trifluoromethyl and methoxy; in particular hydrogen or fluoro; and
30 especially fluoro.

Suitably, X¹ represents halogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy. Typical values of X¹ include fluoro, chloro, methyl, trifluoromethyl and methoxy, especially fluoro.

Typical values of X² include hydrogen and fluoro, especially hydrogen.

In a preferred embodiment, Y represents a chemical bond.

In an alternative embodiment, Y represents a $-OCH_2-$ linkage.

In another embodiment, Y represents an oxygen atom.

In a further embodiment, Y represents a $-NH-$ linkage.

5 Selected values for the substituent Z include phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, furyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrrolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl and tetrazolyl, any of which groups may be optionally substituted by one or more substituents.

10 Particular values of Z include phenyl and triazolyl, either of which groups may be optionally substituted by one or more substituents.

In one favoured embodiment, Z represents an optionally substituted phenyl group, in particular monosubstituted or disubstituted phenyl. In another favoured embodiment, Z represents optionally substituted pyridinyl, especially
15 unsubstituted, monosubstituted or disubstituted pyridin-2-yl, pyridin-3-yl or pyridin-4-yl.

Examples of suitable substituents on the group Z include fluoro, chloro, cyano, nitro, methyl, hydroxy, methoxy, oxy, methanesulphonyl, amino, aminocarbonyl, formyl, methoxycarbonyl and $-CH=NOH$.

20 Examples of typical substituents on the group Z include fluoro, chloro, cyano and methyl.

Examples of particular substituents on the group Z include fluoro and cyano, especially cyano.

Detailed values of Z include cyanophenyl, (cyano)(fluoro)phenyl,
25 (chloro)(cyano)phenyl, nitrophenyl, methoxyphenyl, methanesulphonyl-phenyl, pyridinyl, fluoro-pyridinyl, difluoro-pyridinyl, (amino)(chloro)pyridinyl, cyano-pyridinyl, methyl-pyridinyl, hydroxy-pyridinyl, methoxy-pyridinyl, oxy-pyridinyl, aminocarbonyl-pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, cyano-thienyl, aminocarbonyl-thienyl, formyl-thienyl, methoxycarbonyl-thienyl, thienyl-
30 $CH=NOH$, thiazolyl, isothiazolyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl and methyl-tetrazolyl. Additionally, Z may represent methyl-triazolyl.

Specific values of Z include cyanophenyl, (cyano)(fluoro)phenyl, pyridinyl, difluoro-pyridinyl and cyano-pyridinyl.

Individual values of Z include cyanophenyl, (cyano)(fluoro)phenyl, (chloro)(cyano)phenyl and methyl-triazolyl.

In one embodiment, Z represents cyanophenyl, especially 2-cyanophenyl.

- 5 In another embodiment, Z represents (cyano)(fluoro)phenyl, especially 2-cyano-4-fluorophenyl. Alternative aspects of this embodiment include 2-cyano-5-fluorophenyl, 2-cyano-6-fluorophenyl and 4-cyano-2-fluorophenyl.

In an additional embodiment, Z represents (chloro)(cyano)phenyl, especially 3-chloro-4-cyanophenyl.

- 10 In a further embodiment, Z represents methyl-triazolyl, especially 1-methyl-1*H*[1,2,3]triazol-4-yl or 2-methyl-2*H*[1,2,4]triazol-3-yl.

Typically, R¹ represents hydrocarbon, a heterocyclic group, trifluoromethyl, -COR^a or -CO₂R^a.

- 15 Typical values of R^a include hydrogen and C₁₋₆ alkyl. Suitably, R^a represents hydrogen or methyl.

- Typical values of R^b include hydrogen, C₁₋₆ alkyl, hydroxy(C₁₋₆)alkyl and di(C₁₋₆)alkylamino(C₁₋₆)alkyl. Suitably, R^b represents hydrogen, methyl, ethyl, hydroxyethyl or dimethylaminoethyl. Particular values of R^b include hydrogen, hydroxyethyl and dimethylaminoethyl, especially hydrogen or
20 dimethylaminoethyl.

- Suitable values of R¹ include C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, dihydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, di(C₁₋₆)alkoxy(C₁₋₆)alkyl, cyano(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, C₃₋₇ cycloalkyl, heteroaryl, C₁₋₆ alkyl-heteroaryl, heteroaryl(C₁₋₆)alkyl, trifluoromethyl, formyl, C₂₋₆ alkylcarbonyl and C₂₋₆ alkoxycarbonyl. Additional values of R¹ include trihalo(C₁₋₆)alkyl and (C₁₋₆)alkyl-heteroaryl(C₁₋₆)alkyl.
25

Representative values of R¹ include C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl and trifluoromethyl.

- 30 Particular values of R¹ include C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, trihalo(C₁₋₆)alkyl, heteroaryl(C₁₋₆)alkyl and (C₁₋₆)alkyl-heteroaryl(C₁₋₆)alkyl.

Individual values of R¹ include methyl, ethyl, fluoromethyl, difluoromethyl, hydroxymethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl

(especially 2-hydroxyethyl), fluoroethyl (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), dimethoxyethyl (especially 2,2-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), dihydroxypropyl (especially 1,2-dihydroxyprop-2-yl),
5 fluoropropyl (especially 2-fluoroprop-2-yl), cyanopropyl (especially 2-cyanoprop-2-yl), methoxycarbonylpropyl (especially 2-methoxycarbonylprop-2-yl), *tert* butyl, hydroxybutyl (especially 1-hydroxy-2-methylprop-2-yl), cyclopropyl, pyridinyl, furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, trifluoromethyl, formyl, acetyl and methoxycarbonyl. Additional
10 values of R¹ include trifluoroethyl (especially 2,2,2-trifluoroethyl), methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl) and pyridinylmethyl (especially pyridin-2-ylmethyl).


Specific values of R¹ include methyl, ethyl, fluoroethyl (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), trifluoroethyl (especially
15 2,2,2-trifluoroethyl), isopropyl, methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl) and pyridinylmethyl (especially pyridin-2-ylmethyl).

In a first embodiment, R¹ represents methyl. In a second embodiment, R¹ represents ethyl. In a third embodiment, R¹ represents fluoroethyl (especially 2-fluoroethyl). In a fourth embodiment, R¹ represents difluoroethyl (especially 2,2-
20 difluoroethyl). In a fifth embodiment, R¹ represents trifluoroethyl (especially 2,2,2-trifluoroethyl). In a sixth embodiment, R¹ represents isopropyl. In a seventh embodiment, R¹ represents methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl). In an eighth embodiment, R¹ represents pyridinylmethyl (especially pyridin-2-ylmethyl). In a favoured embodiment, R¹
25 represents 2-hydroxyprop-2-yl. In another embodiment, R¹ represents 2-fluoroprop-2-yl. In an additional embodiment, R¹ represents trifluoromethyl.

In one embodiment individual values of R¹ when -U-V- is -N=CH- or -CH=N- include methyl, ethyl, fluoromethyl, difluoromethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl (especially 2-hydroxyethyl), fluoroethyl
30 (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), dimethoxyethyl (especially 2,2-dimethoxyethyl), isopropyl, hydroxypropyl, dihydroxypropyl (especially 1,2-dihydroxyprop-2-yl), cyanopropyl, methoxycarbonylpropyl (especially 2-methoxycarbonylprop-2-yl), *tert* butyl, hydroxybutyl (especially 1-hydroxy-2-methylprop-2-yl), cyclopropyl, pyridinyl,

furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, trifluoromethyl, formyl, acetyl and methoxycarbonyl.

A particular sub-class of compounds according to the invention is represented by the compounds of formula IIA, and pharmaceutically acceptable salts thereof:



(IIA)

wherein

10 U, V and Z are as defined above;

X¹¹ represents hydrogen, fluoro, chloro, methyl, trifluoromethyl or methoxy;

X¹² represents hydrogen or fluoro; and

R¹¹ represents C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, trihalo(C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, dihydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, di(C₁₋₆)alkoxy(C₁₋₆)alkyl, cyano(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, C₃₋₇ cycloalkyl, heteroaryl, C₁₋₆ alkyl-heteroaryl, heteroaryl(C₁₋₆)alkyl, (C₁₋₆)alkyl-heteroaryl(C₁₋₆)alkyl, trifluoromethyl, formyl, C₂₋₆ alkylcarbonyl or C₂₋₆ alkoxycarbonyl.

20 The present invention also provides a compound of formula IIA as depicted above, or a pharmaceutically acceptable salt thereof, wherein

-U-V- represents -CH=CH-;

R¹¹ represents C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, dihydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, di(C₁₋₆)alkoxy(C₁₋₆)alkyl, cyano(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, C₃₋₇ cycloalkyl, heteroaryl, C₁₋₆ alkyl-heteroaryl, heteroaryl(C₁₋₆)alkyl, trifluoromethyl, formyl, C₂₋₆ alkylcarbonyl or C₂₋₆ alkoxycarbonyl; and

Z, X¹¹ and X¹² are as defined above.

Suitable values of X¹¹ include hydrogen and fluoro, especially fluoro.

Typical values of X¹¹ include fluoro, chloro, methyl, trifluoromethyl and methoxy.

5 A particular value of X¹¹ is fluoro.

In a favoured embodiment, X¹² represents hydrogen. In another embodiment, X¹² represents fluoro.

Where R¹¹ represents heteroaryl, this group is suitably pyridinyl, furyl, thienyl or oxazolyl.

10 Where R¹¹ represents C₁₋₆ alkyl-heteroaryl, this group is suitably methylthiazolyl (e.g. 2-methylthiazol-5-yl) or methyloxadiazolyl (e.g. 3-methyl-[1,2,4]oxadiazol-5-yl).

Where R¹¹ represents heteroaryl(C₁₋₆)alkyl, this group is suitably imidazolylmethyl or triazolylmethyl. Additionally, the

15 heteroaryl(C₁₋₆)alkyl group R¹¹ may be pyridinylmethyl.

Representative values of R¹¹ include C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl and trifluoromethyl.

Particular values of R¹¹ include C₁₋₆ alkyl, halo(C₁₋₆)alkyl, dihalo(C₁₋₆)alkyl, trihalo(C₁₋₆)alkyl, heteroaryl(C₁₋₆)alkyl and (C₁₋₆)alkyl-
20 heteroaryl(C₁₋₆)alkyl.


Individual values of R¹¹ include methyl, ethyl, fluoromethyl, difluoromethyl, hydroxymethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl (especially 2-hydroxyethyl), fluoroethyl (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), dimethoxyethyl (especially 2,2-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), dihydroxypropyl (especially 1,2-dihydroxyprop-2-yl), fluoropropyl (especially 2-fluoroprop-2-yl), cyanopropyl (especially 2-cyanoprop-2-yl), methoxycarbonylpropyl (especially 2-methoxycarbonylprop-2-yl), *tert*-butyl, hydroxybutyl (especially 1-hydroxy-2-methylprop-2-yl), cyclopropyl, pyridinyl, 25 furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, trifluoromethyl, formyl, acetyl and methoxycarbonyl. Additional values of R¹¹ include trifluoroethyl (especially 2,2,2-trifluoroethyl), methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl) and 30 pyridinylmethyl (especially pyridin-2-ylmethyl).

Specific values of R¹¹ include methyl, ethyl, fluoroethyl (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), trifluoroethyl (especially 2,2,2-trifluoroethyl), isopropyl, methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl) and pyridinylmethyl (especially pyridin-2-ylmethyl).

5 In a first embodiment, R¹¹ represents methyl. In a second embodiment, R¹¹ represents ethyl. In a third embodiment, R¹¹ represents fluoroethyl (especially 2-fluoroethyl). In a fourth embodiment, R¹¹ represents difluoroethyl (especially 2,2-difluoroethyl). In a fifth embodiment, R¹¹ represents trifluoroethyl (especially 2,2,2-trifluoroethyl). In a sixth embodiment, R¹¹ represents isopropyl.
10 In a seventh embodiment, R¹¹ represents methyl-triazolylmethyl (especially 2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl). In an eighth embodiment, R¹¹ represents pyridinylmethyl (especially pyridin-2-ylmethyl). In a favoured embodiment, R¹¹ represents 2-hydroxyprop-2-yl. In another embodiment, R¹¹ represents 2-fluoroprop-2-yl. In an additional embodiment, R¹¹ represents trifluoromethyl.

15 When -U-V- is -N=CH- or -CH=N- individual values of R¹¹ include methyl, ethyl, fluoromethyl, difluoromethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl (especially 2-hydroxyethyl), fluoroethyl (especially 2-fluoroethyl), difluoroethyl (especially 2,2-difluoroethyl), dimethoxyethyl (especially 2,2-dimethoxyethyl), isopropyl, hydroxypropyl, dihydroxypropyl (especially 1,2-
20 dihydroxyprop-2-yl), cyanopropyl, methoxycarbonylpropyl (especially 2-methoxycarbonylprop-2-yl), *tert*-butyl, hydroxybutyl (especially 1-hydroxy-2-methylprop-2-yl), cyclopropyl, pyridinyl, furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, trifluoromethyl, formyl, acetyl and methoxycarbonyl.

25 One representative subset of the compounds of formula IIA above is represented by the compounds of formula IIB, and pharmaceutically acceptable salts thereof:



(IIB)


wherein U, V, X¹¹, X¹² and R¹¹ are as defined above; and

R² represents hydrogen or fluoro.

5 In one embodiment, R² is hydrogen.

In another embodiment, R² is fluoro, in which case the fluorine atom R² is favourably attached to the phenyl ring at the 4-, 5- or 6-position (relative to the cyano group at position 2).

10 Another representative subset of the compounds of formula IIA above is represented by the compounds of formula IIC, and pharmaceutically acceptable salts thereof:



(IIC)

15 wherein U, V, X¹¹, X¹² and R¹¹ are as defined above; and

R³ represents hydrogen, fluoro, cyano or methyl.


In one embodiment, R³ is hydrogen.

In an additional embodiment, R³ is fluoro.

In another embodiment, R³ is cyano.

In a further embodiment, R³ is methyl.

A further representative subset of the compounds of formula IIA above is
 5 represented by the compounds of formula IID, and pharmaceutically acceptable salts thereof:



(IID)

10 wherein U, V, X¹¹, X¹², R³ and R¹¹ are as defined above; and

R⁴ represents hydrogen or fluoro.

Suitably, R⁴ represents hydrogen.

In another embodiment, R⁴ represents fluoro.

The present invention also provides a compound of formula IIB, IIC or IID
 15 as depicted above, or a pharmaceutically acceptable salt thereof, wherein

-U-V- represents -CH=CH-; and

X¹¹, X¹², R², R³, R⁴ and R¹¹ are as defined above.

Specific compounds within the scope of the present invention include:

2'-fluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-a]pyrazin-3-yl)-biphenyl-2-
 20 carbonitrile;

5'-(7-ethyl-8-oxo-7,8-dihydroimidazo[1,2-a]pyrazin-3-yl)-2'-fluorobiphenyl-2-
 carbonitrile;

2'-fluoro-5'-[7-(2-fluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-a]pyrazin-3-
 yl]biphenyl-2-carbonitrile;

25 5'-[7-(2,2-difluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-a]pyrazin-3-yl]-2'-
 fluorobiphenyl-2-carbonitrile;

- 4,2'-difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
- 4,2'-difluoro-5'-(7-ethyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
- 5 4,2'-difluoro-5'-[7-(2,2-difluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 4,2'-difluoro-5'-[7-(1-methylethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 3-chloro-2'-fluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-4-carbonitrile;
- 10 4,2'-difluoro-5'-[7-(2-fluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 4'-fluoro-3'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
- 15 5'-fluoro-3'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
- 2,2'-difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-4-carbonitrile;
- 4,2'-difluoro-5'-[7-(2,2,2-trifluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 20 4,2'-difluoro-5'-[7-(2-methyl-2*H*[1,2,4]triazol-3-ylmethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 3-[4-fluoro-3-(1-methyl-1*H*[1,2,3]triazol-4-ylmethoxy)phenyl]-7-methyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one;
- 25 3-[4-fluoro-3-(2-methyl-2*H*[1,2,4]triazol-3-ylmethoxy)phenyl]-7-methyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one;
- 7-ethyl-3-[4-fluoro-3-(2-methyl-2*H*[1,2,4]triazol-3-ylmethoxy)phenyl]-7*H*-imidazo[1,2-*a*]pyrazin-8-one;
- 4,2'-difluoro-5'-[8-oxo-7-(pyridin-2-ylmethyl)-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile;
- 30 4,2'-difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
- 4,2'-difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;

- 5,2'-difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
6,2'-difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
5 6,2'-difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
6,2'-difluoro-5'-(7-ethyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile;
5,2'-difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
10 4,2'-difluoro-5'-[8-oxo-7-(pyridin-2-ylmethyl)-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
6,2'-difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile;
15 and pharmaceutically acceptable salts thereof.

Also provided is:

- 4,2'-difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*d*][1,2,4]triazin-3-yl)biphenyl-2-carbonitrile;
and pharmaceutically acceptable salts thereof.

- 20 Also provided is:

3'-(1-methyl-6-oxo-1,6-dihydropurin-9-yl)biphenyl-2-carbonitrile;
and pharmaceutically acceptable salts thereof.

- Also provided by the present invention is a method for the treatment and/or prevention of anxiety which comprises administering to a patient in need
25 of such treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

- Further provided by the present invention is a method for the treatment and/or prevention of convulsions (e.g. in a patient suffering from epilepsy or a related disorder) which comprises administering to a patient in need of such
30 treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

Details of the binding affinity of the claimed compounds, their potentiation of the GABA EC₂₀ response, measurement of anxiolytic and anticonvulsant

activity, and pharmaceutical compositions can be found on pages 27-30 of WO-A-02074773.

In another aspect, the present invention provides a method for the treatment and/or prevention of cognitive disorders, including dementing
5 conditions such as Alzheimer's disease, which comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.


Cognitive disorders for which the compounds of the present invention may be of benefit include delirium, dementia, amnestic disorders, and cognition
10 deficits, including age-related memory deficits, due to traumatic injury, stroke, Parkinson's disease and Down Syndrome. Any of these conditions may be attributable to substance abuse or withdrawal. Examples of dementia include dementia of the Alzheimer's type with early or late onset, and vascular dementia, any of which may be uncomplicated or accompanied by delirium, delusions or
15 depressed mood; and dementia due to HIV disease, head trauma, Parkinson's disease or Creutzfeld-Jakob disease.

In order to elicit their behavioural effects, the compounds of the invention will ideally be brain-penetrant; in other words, these compounds will be capable of crossing the so-called "blood-brain barrier". Preferably, the compounds of the
20 invention will be capable of exerting their beneficial therapeutic action following administration by the oral route.

Unit dosage forms comprising the claimed compounds contain from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg,
25 of the active ingredient.

In the treatment of neurological disorders, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day.

30 The compounds in accordance with the present invention may be prepared by a process which comprises attachment of the R¹ moiety to a compound of formula III:




wherein U, V, X¹, X², Y and Z are as defined above; by conventional alkylation or acylation methods.

- 5 For instance, where R¹ in the compounds of formula I above represents an optionally substituted C₁₋₆ alkyl group, the moiety R¹ may be attached by treating the appropriate compound of formula III with a suitable alkyl halide, e.g. iodomethane, iodoethane, 1-bromo-2-fluoroethane, 2-bromo-1,1-difluoroethane, 1-iodo-2,2,2-trifluoroethane or 2-bromopropane, typically in the presence of a base
- 10 such as sodium hydride. Similarly, where R¹ represents an optionally substituted heteroaryl(C₁₋₆)alkyl group, the moiety R¹ may be attached by reacting the appropriate compound of formula III with a suitable alkylating agent, e.g. 3-chloromethyl-2-methyl-2H-[1,2,4]triazole or 2-chloromethylpyridine, typically in the presence of a base such as sodium hydride. Alternatively, where R¹ in the
- 15 compounds of formula I above represents methyl, the methyl group R¹ may be attached by treating the appropriate compound of formula III with a strong base such as hexamethyldisilazane, followed by chloro(chloromethyl)dimethylsilane; and subsequently treating the compound thereby obtained with cesium fluoride.


20 Except where X¹ and X² both simultaneously represent hydrogen, the compounds of formula III above are novel compounds and represent a further feature of the present invention.

The compounds of formula III may suitably be prepared from the appropriate methoxy-substituted compound of formula IV:




wherein U, V, X¹, X², Y and Z are as defined above; by treatment with hydrogen bromide, typically in acetic acid.

- 5 In another procedure, the compounds in accordance with the present invention may be prepared by a process which comprises reacting a compound of formula V with a compound of formula VI:



(V)



(VI)

10

wherein U, V, X¹, X², Y, Z and R¹ are as defined above, L¹ represents a suitable leaving group, and M¹ represents a boronic acid moiety -B(OH)₂ or a cyclic ester thereof formed with an organic diol, e.g. pinacol, 1,3-propanediol or neopentyl glycol, or M¹ represents -Sn(Alk)₃ in which Alk represents a C₁₋₆ alkyl group,


15 typically *n*-butyl; in the presence of a transition metal catalyst.

The leaving group L¹ is typically a halogen atom, e.g. bromo.

The transition metal catalyst of use in the reaction between compounds V and VI is suitably palladium(II) acetate, in which case the reaction is typically accomplished in the presence of triphenylphosphine. The reaction is conveniently

20 carried out at an elevated temperature in a solvent such as 1,4-dioxane, typically in the presence of sodium carbonate.

The intermediates of formula V may be prepared by attaching the R¹ moiety to a compound of formula VII:




(VII)

5

wherein U, V and L¹ are as defined above; under conditions analogous to those described above for attachment of the R¹ moiety to a compound of formula III.

The intermediates of formula VII may be prepared from the appropriate methoxy-substituted precursor of formula VIII:




(VIII)

10


wherein U, V and L¹ are as defined above; by treatment with hydrogen bromide, typically in acetic acid.

15

In a further procedure, the compounds in accordance with the present invention may be prepared by a process which comprises reacting a compound of formula IX with a compound of formula X:



(IX)




(X)

wherein U, V, X¹, X², Y, Z, R¹ and L¹ are as defined above; in the presence of a transition metal catalyst.

5 The transition metal catalyst of use in the reaction between compounds IX and X is suitably palladium(II) acetate, in which case the reaction is typically accomplished in the presence of triphenylphosphine. The reaction is conveniently carried out at an elevated temperature in a solvent such as *N,N*-dimethylacetamide, typically in the presence of potassium acetate.

10 The intermediates of formula IX where -U-V- is -N=C- may suitably be prepared by attachment of the R¹ moiety to the compound of formula XI:




(XI)

by conventional alkylation or acylation methods.

15 For instance, where R¹ in the compounds of formula I above represents an optionally substituted C₁₋₆ alkyl group, the moiety R¹ may be attached by treating the appropriate compound of formula IX with a suitable alkyl halide, e.g. iodomethane, iodoethane, 1-bromo-2-fluoroethane or 2-bromo-1,1-difluoroethane, typically in the presence of a base such as sodium hydride. Alternatively, where
20 R¹ in the compounds of formula I above represents methyl, the methyl group R¹ may be attached by treating the appropriate compound of formula IX with a strong base such as hexamethyldisilazane, followed by chloro(chloromethyl)-dimethylsilane; and subsequently treating the compound thereby obtained with cesium fluoride.

25 The intermediate of formula XI may be prepared by reacting the compound of formula XII:




(XII)

with *N,N*-dimethylformamide dimethyl acetal, typically by heating in Dowtherm A.

5 The intermediate of formula XII may be obtained from the process described in EP-A-0713876.

The compounds in accordance with the present invention particularly those in which -U-V- is -CH=N-, may alternatively be prepared by a process which comprises reacting a compound of formula XIII:

10




(XIII)

wherein U, V, X¹, X², Y, Z and R¹ are as defined above; with formic acid.

15 The reaction is conveniently accomplished by stirring the reactants at an elevated temperature, typically at the reflux temperature of the reaction mixture.

The intermediates of formula XIII may be prepared by reduction of a compound of formula XIV:




(XIV)


wherein U, V, X¹, X², Y, Z and R¹ are as defined above.

The reduction is conveniently effected by treating compound XIV with a
 5 reducing agent, e.g. tin(II) chloride, typically at an elevated temperature in a
 suitable solvent, for example a mixture of ethanol and tetrahydrofuran.

The intermediates of formula XIV may be prepared by reacting a
 compound of formula XV with a compound of formula XVI:



(XV)




(XVI)

wherein U, V, X¹, X², Y, Z and R¹ are as defined above, and L¹ represents a
 suitable leaving group.


The leaving group L¹ is typically a halogen atom, e.g. chloro.

15 The reaction between compounds XV and XVI is conveniently effected by
 heating the reactants, to a temperature typically in the region of 90°C, in the
 presence of a base such as triethylamine, in a solvent such as dimethylsulfoxide.

In another procedure, the compounds according to the present invention,
 particularly those in which -U-V- is -CH=N-, in which Y represents a chemical
 20 bond may be prepared by a process which comprises reacting a compound of
 formula XVII with a compound of formula XVIII:



(XVII)




(XVIII)

wherein U, V, X¹, X², Z and R¹ are as defined above, L² represents a suitable
 5 leaving group, and M¹ represents a boronic acid moiety -B(OH)₂ or a cyclic ester
 thereof formed with an organic diol, e.g. pinacol, 1,3-propanediol or neopentyl
 glycol, or M¹ represents -Sn(Alk)₃ in which Alk represents a C₁₋₆ alkyl group,
 typically *n*-butyl; in the presence of a transition metal catalyst.


10 In the compounds of formula XVII above, the leaving group L² is typically
 trifluoromethanesulfonyloxy (triflyloxy); or a halogen atom, e.g. bromo.

The transition metal catalyst of use in the reaction between compounds
 XVII and XVIII is suitably tetrakis(triphenylphosphine)-palladium(0). The
 reaction is conveniently carried out at an elevated temperature in a solvent such
 as *N,N*-dimethylacetamide, 1,2-dimethoxyethane, 1,4-dioxane or tetrahydrofuran,
 15 advantageously in the presence of potassium phosphate, copper(I) iodide, sodium
 carbonate or cesium carbonate. Alternatively, the transition metal catalyst
 employed may be tris(dibenzylideneacetone)dipalladium(0), in which case the
 reaction is conveniently effected at an elevated temperature in a solvent such as
 1,4-dioxane, typically in the presence of tri-*tert*-butylphosphine and potassium
 20 phosphate.

Alternatively, the compounds according to the present invention,
 particularly those in which -U-V- is -CH=N-, in which Y represents a chemical
 bond may be prepared by a process which comprises reacting a compound of
 formula XIX with a compound of formula XX:




(XIX)



(XX)

wherein U, V, X¹, X², Z, R¹, L² and M¹ are as defined above; in the presence of a transition metal catalyst; under conditions analogous to those described above for the reaction between compounds XVII and XVIII.

In an additional procedure, the compounds according to the present invention, particularly those in which -U-V- is -CH=N-, in which Y represents an oxygen atom may be prepared by a process which comprises reacting a compound of formula XX as defined above with a compound of formula XXI:




(XXI)

wherein U, V, X¹, X² and R¹ are as defined above.

The reaction is conveniently carried out under basic conditions, e.g. using sodium hydride in a solvent such as *N,N*-dimethylformamide, typically at an elevated temperature which may be in the region of 120°C.

In a further procedure, the compounds according to the present invention, particularly those in which -U-V- is -CH=N-, in which Y represents a -NH-

linkage may be prepared by a process which comprises reacting a compound of formula XX as defined above with a compound of formula XXII:



(XXII)

5

wherein U, V, X¹, X² and R¹ are as defined above.


In relation to the reaction between compounds XX and XXII, the leaving group L² in the compounds of formula XX may suitably represent fluoro.

The reaction between compounds XX and XXII is conveniently carried out by heating the reactants, typically at a temperature in the region of 120°C, in a solvent such as *N,N*-dimethylformamide.

Where M¹ in the intermediates of formula XVIII and XIX above represents a boronic acid moiety -B(OH)₂ or a cyclic ester thereof formed with pinacol or neopentyl glycol, the relevant compound XVIII or XIX may be prepared by reacting bis(pinacolato)diboron or bis(neopentyl glycolato)diborane respectively with a compound of formula XX or XVII as defined above; in the presence of a transition metal catalyst.

The transition metal catalyst of use in the reaction between bis(pinacolato)diboron or bis(neopentyl glycolato)diborane and compound XX or XVII is suitably dichloro[1,1'-bis(diphenylphosphino)ferrocene]-palladium(II). The reaction is conveniently carried out at an elevated temperature in a solvent such as 1,4-dioxane, optionally in admixture with dimethylsulfoxide, typically in the presence of 1,1'-bis(diphenylphosphino)ferrocene and/or potassium acetate.


The intermediates of formula XVII above may be prepared by reacting a compound of formula XXIII:



(XXIII)

wherein U, V, X¹, X², R¹ and L² are as defined above; with formic acid; under conditions analogous to those described above for the reaction between compound
 5 XIII and formic acid.

The intermediates of formula XXIII may be prepared by reaction of a compound of formula V as defined above with a compound of formula XXIV:



(XXIV)


10

wherein X¹, X² and L² are as defined above; under conditions analogous to those described above for the reaction between compounds XV and XVI; followed by reduction of the nitro group in the resulting compound, under conditions analogous to those described above for the reduction of the nitro group in
 15 compound XIV.

Where L² in the intermediates of formula XVII above represents triflyloxy, the relevant compound XVII may be prepared by reacting the appropriate compound of formula XXI as defined above with triflic anhydride, typically in the presence of pyridine.

20

The intermediates of formula XXI above may suitably be prepared from the appropriate methoxy-substituted precursor of formula XXV:




(XXV)

wherein U, V, X¹, X² and R¹ are as defined above; by treatment with boron
 5 tribromide, typically in chloroform or dichloromethane; or with hydrogen
 bromide, typically in acetic acid at reflux.

The intermediates of formula XXII and XXV above may be prepared by
 reacting a compound of formula XV as defined above with the appropriate
 compound of formula XXVI:

10



(XXVI)

wherein X¹ and X² are as defined above, and Y¹ represents amino or methoxy;
 under conditions analogous to those described above for the reaction between
 15 compounds XV and XVI; followed by reduction of the nitro group in the resulting
 compound, under conditions analogous to those described above for the reduction
 of the nitro group in compound XIV; followed by reaction of the product thereby
 obtained with formic acid, under conditions analogous to those described above
 for the reaction between compound XIII and formic acid.

20

The compounds according to the invention in which Y represents a
 chemical bond and Z represents pyrrol-1-yl may be prepared by reacting a

compound of formula XXII as defined above with 2,5-dimethoxy-tetrahydrofuran. The reaction is conveniently accomplished at an elevated temperature in a solvent such as acetic acid.

The intermediates of formula III, IV, V, VII, VIII and IX wherein
5 -U-V- represents -CH₂-CH₂- may be prepared from the corresponding compound wherein -U-V- represents -CH=CH- by catalytic hydrogenation.

The compounds of formula IV, VI, VIII, IX and X may conveniently be prepared by the procedures described in WO 02/10170, or by methods analogous thereto.

10 Where they are not commercially available, the starting materials of formula XV, XVI, XX, XXIV and XXVI may be prepared by methods analogous to those described in the accompanying Examples, or by standard methods well known from the art.

It will be understood that any compound of formula I initially obtained
15 from any of the above processes may, where appropriate, subsequently be elaborated into a further compound of formula I by techniques known from the art and detailed on pages 40-42 of WO-A-02074773. A compound of formula I wherein -U-V- represents -CH=CH- may be converted into the corresponding compound wherein

20 -U-V- represents -CH₂-CH₂- by catalytic hydrogenation.

Mixtures of products may be purified and intermediates protected as described on pages 42-43 of WO-A-02074773.

The compounds of the accompanying Examples were tested in the assay described on pages 43 and 44 of WO-A-02074773, and all were found to possess a
25 K_i value for displacement of [³H]-flumazenil from the α₂ and/or α₃ and/or α₅ subunit of the human GABA_A receptor of 100 nM or less.

The following Examples illustrate the preparation of compounds according to the invention.

30 **EXAMPLE 1** 2'-Fluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-
a]pyrazin-3-yl)-biphenyl-2-carbonitrile

A suspension of 2,3-dichloropyrazine (35 g, 0.23 mol) in 25% aqueous ammonia (200 ml) was heated at 100°C for 12 h in a PTFE-lined pressure reactor (terminal pressure 100 psi). The reaction was cooled to ambient temperature and

the resulting crystalline solid collected by filtration. This solid was triturated with water (150 ml) and dried to afford 2-amino-3-chloropyrazine as a buff-coloured crystalline solid (28 g, 92%): δ_{H} (400 MHz, DMSO) 6.78 (2H, br s), 7.56 (1H, d, J 2.5), 7.95 (1H, d, J 2.5).

5 Bromoacetaldehyde diethyl acetal (45 ml, 0.29 mol) was treated with water (33 ml) and 48% hydrobromic acid (33 ml) and this mixture was heated at 95°C for 90 min. The reaction was cooled, diluted with propan-2-ol (300 ml) and treated with sodium hydrogencarbonate (33 g) added in portions. This mixture was stirred for 30 min then filtered. The filtrate was treated with 2-amino-3-
10 chloropyrazine (25 g, 0.19 mol) and then heated at 90°C for 16 h. The reaction was cooled to ambient temperature, concentrated to about one-third volume and then treated with 48% hydrobromic acid (25 ml). More propan-2-ol (300 ml) was added and the mixture aged for 1 h. The resulting solid was collected by filtration, washed with propan-2-ol and then dissolved in water (500 ml). This
15 solution was made basic by adding solid sodium hydrogencarbonate and then extracted with chloroform (3 x 250 ml). The organics were combined, dried over anhydrous magnesium sulphate, filtered and concentrated to give a solid. Trituration with diethyl ether afforded 8-chloroimidazo[1,2-*a*]pyrazine as an off-white solid (18.6 g, 63%): δ_{H} (360 MHz, DMSO) 7.73 (1H, d, J 4.5), 7.87 (1H, d, J
20 1), 8.28 (1H, d, J 1), 8.67 (1H, d, J 4.5).

A cooled (0°C) suspension of 8-chloroimidazo[1,2-*a*]pyrazine (18.6 g, 0.12 mol) and sodium acetate (29.8 g, 0.36 mol) in methanol (125 ml, pre-saturated with solid potassium bromide) was treated with bromine (6.5 ml, 0.13 mol) added dropwise over 5 min. After stirring for 10 min thin-layer chromatography
25 indicated no starting material. Solid sodium sulphite (15.3 g, 0.12 mol) was then added to the slurry and stirring continued for 10 min. The mixture was then treated with saturated aqueous sodium hydrogencarbonate (650 ml) added in portions. This mixture was extracted with dichloromethane (2 x 350 ml). The organics were combined, dried over anhydrous magnesium sulphate, filtered and
30 concentrated to afford 3-bromo-8-chloroimidazo[1,2-*a*]pyrazine as a cream-coloured solid: δ_{H} (400 MHz, CDCl_3) 7.82 (1H, d, J 4.5), 7.83 (1H, s), 8.05 (1H, d, J 4.5). This solid was suspended in 1:1 dichloromethane/methanol (150 ml) and treated with solid sodium methoxide (9.8 g, 0.18 mol). The resulting mixture was then stirred at 40°C for 2 h. The reaction was cooled, diluted with water (750 ml)

then extracted with dichloromethane (600 ml). The organics were dried over anhydrous magnesium sulphate, filtered and concentrated to give 3-bromo-8-methoxyimidazo[1,2-*a*]pyrazine as a white solid (24.5 g, 89% over 2 steps): δ_{H} (360 MHz, DMSO) 4.06 (3H, s), 7.57 (1H, d, *J* 4.5), 7.81 (1H, s), 8.03 (1H, d, *J* 4.5).

5 A mixture of the above product (680 mg, 3 mmol) and 2'-fluoro-5'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-biphenyl-2-carbonitrile (prepared as described in WO 02/074773) (1.45 g, 4.5 mmol) in tetrahydrofuran (9 ml) was treated with 2M sodium carbonate (3 ml) then degassed with nitrogen for 10 min. Tetrakis(triphenylphosphine)palladium(0) (100 mg, 0.09 mmol) was added and
10 this mixture was heated under reflux for 12 h. The reaction was cooled and the majority of the solvent removed on a rotary evaporator. The residue was partitioned between dichloromethane and water. The organics were washed with water, brine, dried over anhydrous magnesium sulphate, filtered and concentrated *in vacuo*. Purification of the residue by chromatography on silica
15 gel eluting with 2% methanol in dichloromethane gave 2'-fluoro-5'-(8-methoxyimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile as a cream solid (768 mg, 74%): δ_{H} (400 MHz, CDCl_3) 4.19 (3H, s), 7.38-7.46 (2H, m), 7.53-7.64 (4H, m), 7.71 (1H, td, *J* 8 and 1.5), 7.73 (1H, s), 7.85 (1H, dd, *J* 8 and 1), 8.02 (1H, d, *J* 4.5); m/z (ES^+) 345 [$\text{M}+\text{H}$] $^+$.

20 A suspension of the above product (750 mg, 2.2 mmol) in hydrogen bromide (30 wt % in acetic acid, 10 ml) was heated at 95°C for 45 min. The reaction was cooled, poured into ice water and neutralised by the addition of solid sodium hydrogencarbonate. The resulting solid was collected by filtration and air-dried, then triturated with ether, to afford 2'-fluoro-5'-(8-oxo-7,8-
25 dihydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile as a cream-coloured solid (550 mg, 76%): δ_{H} (360 MHz, DMSO) 6.92 (1H, t, *J* 6), 7.49 (1H, d, *J* 5), 7.60-7.71 (2H, m), 7.75-7.89 (5H, m), 8.03 (1H, dd, *J* 7 and 0.5), 11.38 (1H, d, *J* 5.5); m/z (ES^+) 331 [$\text{M}+\text{H}$] $^+$.

30 A suspension of the above product (250 mg, 0.75 mmol) in acetonitrile (10 ml) was treated with 1,1,1,3,3,3-hexamethyldisilazane (90 μl , 0.41 mmol) and then heated at reflux for 2 h. The resulting gel-like mixture was diluted with acetonitrile (5 ml) then treated with chloro(chloromethyl)-dimethylsilane (110 μl , 0.83 mmol) and heating at reflux continued for 24 h. The reaction was cooled, the acetonitrile removed *in vacuo* and the residue treated with 1,2-dimethoxyethane

(12 ml). Caesium fluoride (160 mg, 1.4 mmol) was then added and the reaction heated at reflux for 6 h. After cooling to ambient temperature the mixture was dissolved in 1:1 dichloromethane/methanol and pre-adsorbed onto silica.

Purification by chromatography on silica eluting with 4% methanol in

- 5 dichloromethane (containing 0.5% ammonia) followed by 8% methanol in dichloromethane (containing 0.5% ammonia) gave the title compound (65 mg, 25%): δ_{H} (400 MHz, DMSO) 3.47 (3H, s), 7.19 (1H, d, J 6), 7.57-7.88 (8H, m), 8.04 (1H, dd, J 8 and 1); m/z (ES⁺) 345 [M+H]⁺.

10 **EXAMPLE 2** 5'-(7-Ethyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-2'-fluorobiphenyl-2-carbonitrile

- A suspension of 2'-fluoro-5'-(8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile (50.0 mg, 0.51 mmol) in 1,2-dimethoxyethane (2 ml) and *N,N*-dimethylformamide (0.5 ml) was treated with sodium hydride (6.4 mg of
15 a 60% dispersion in mineral oil, 0.16 mmol). After stirring at ambient temperature for 10 min, lithium bromide (26.3 mg, 0.30 mmol) was added, and stirring continued for 15 min. Iodoethane (24.2 μ l, 0.30 mmol) was added and the solution heated at 65°C for 18 h. Water (15 ml) was added and the resulting mixture was extracted with dichloromethane (3 x 10 ml). The combined organic
20 layers were washed with water (10 ml), saturated sodium chloride solution (10 ml), dried over anhydrous magnesium sulphate then concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel, eluting with 2% methanol in dichloromethane. Crystallisation from methanol afforded the title
25 compound as a white solid (23.9 mg, 44%): δ_{H} (500 MHz, CDCl₃) 1.38 (3H, t, J 7.2), 4.06 (2H, q, J 7.2), 6.72 (1H, d, J 5.9), 7.39 (1H, t, J 9.3), 7.44 (1H, d, J 5.9), 7.53-7.61 (5H, m), 7.69-7.73 (1H, m), 7.84 (1H, d, J 7.3); m/z (ES⁺) 359 [M+H]⁺.

Examples 3 and 4 are made by the method of Example 2 using 1-bromo-2-fluoroethane and 2-bromo-1,1-difluoroethane respectively.

EXAMPLE 3 2'-Fluoro-5'-[7-(2-fluoroethyl)-8-oxo-7,8-dihydroimidazo
[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

δ_H (500 MHz, $CDCl_3$) 4.27 (1H, t, J 4.5), 4.32 (1H, t, J 4.5), 4.71 (1H, t, J 4.4), 4.81 (1H, t, J 4.5), 6.80 (1H, d, J 5.9), 7.40-7.43 (2H, m), 7.53-7.61 (4H, m),
5 7.63 (1H, s), 7.69-7.73 (1H, m), 7.85 (1H, d, J 7.7); m/z (ES^+) 377 $[M+H]^+$.

EXAMPLE 4 5'-[7-(2,2-Difluoroethyl)-8-oxo-7,8-dihydroimidazo
[1,2-*a*]pyrazin-3-yl]-2'-fluorobiphenyl-2-carbonitrile

δ_H (500 MHz, $CDCl_3$) 4.31 (2H, td, J 13 and 4.4), 6.14 (1H, tt, J 56 and 4.4),
10 6.74 (1H, d, J 6.1), 7.41 (1H, t, J 9.3), 7.46 (1H, d, J 6.1), 7.55-7.60 (4H, m), 7.65
(1H, s), 7.70-7.73 (1H, m), 7.85 (1H, d, J 7.1); m/z (ES^+) 395 $[M+H]^+$.

EXAMPLE 5 4,2'-Difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo
[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

15 A suspension of 3-bromo-8-methoxyimidazo[1,2-*a*]pyrazine (1.6 g, 7 mmol)
in hydrogen bromide (30 wt % in acetic acid, 15 ml) was heated at 80°C for 90
min. The reaction was cooled, diluted with water (75 ml) then neutralised with
solid sodium hydrogencarbonate. The resulting solid was collected by filtration,
washed with water then dried under vacuum to afford 3-bromo-7*H*-imidazo[1,2-
20 *a*]pyrazin-8-one as a white powder: δ_H (360 MHz, DMSO) 6.99 (1H, d, J 5.6), 7.29
(1H, d, J 5.6), 7.62 (1H, s). This powder was suspended in *N,N*-
dimethylformamide (15 ml) then treated with sodium hydride (203 mg of a 60%
dispersion in mineral oil, 8.4 mmol). The resulting mixture was heated at 60°C
for 20 min then treated with iodomethane (870 μ l, 14 mmol). After stirring for 15
25 min the reaction was cooled, diluted cautiously with water (150 ml) and then
extracted with dichloromethane (2 x 100 ml). The organics were combined,
washed with water, dried over anhydrous magnesium sulphate, filtered and
concentrated under high vacuum. The residue was triturated with diethyl ether
and the resulting solid collected by filtration to furnish 3-bromo-7-methyl-7*H*-
30 imidazo[1,2-*a*]pyrazin-8-one as a white solid (1.4 g, 88% over 2 steps): δ_H (360
MHz, DMSO) 3.45 (3H, s), 7.28 (1H, d, J 6), 7.39 (1H, d, J 5.6), 7.62 (1H, s).

A mixture of the above product (171 mg, 0.75 mmol), 4,2'-difluoro-5'-(5,5-
dimethyl-[1,3,2]dioxaborinan-2-yl)biphenyl-2-carbonitrile (prepared as described
in WO 02/074773) (392 mg, 1.2 mmol), palladium(II) acetate (7 mg, 0.03 mmol)

and triphenylphosphine (8 mg, 0.03 mmol) in 1,4-dioxane (3 ml) and 2M aqueous sodium carbonate (0.75 ml) was heated at 80°C for 3 h. The reaction was cooled then diluted with ethyl acetate (25 ml). The mixture was extracted with 4N hydrochloric acid (25 ml) and the organics discarded. The aqueous was washed
5 with ethyl acetate, filtered through GF/A glass microfibre filter paper and the filtrate made basic with solid sodium hydrogencarbonate. The resulting solid was collected by filtration, triturated with water and dried under high vacuum to afford the title compound as a cream-coloured solid (175 mg, 64%): δ_{H} (500 MHz, DMSO) 3.47 (3H, s), 7.19 (1H, d, J 6.0), 7.57-7.61 (2H, m), 7.72 (1H, s), 7.76-7.86
10 (4H, m), 8.08 (1H, dd, J 9 and 3); m/z (ES⁺) 363 [M+H]⁺.

Examples 6 to 11 were made by the method of Example 5.

EXAMPLE 6 4'-Fluoro-3'-(7-methyl-8-oxo-7,8-dihydroimidazo
15 [1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

4'-Fluoro-3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile was coupled to give the title compound as an off-white solid (40 mg, 24%): δ_{H} (400 MHz, d₆-DMSO) 3.48 (3H, s), 7.20 (1H, d, J 6), 7.43 (1H, dd, J 6 and 2), 7.61-7.66 (2H, m), 7.72-7.86 (5H, m), 8.01 (1H, dd, J 8 and 1); m/z (ES⁺) 345
20 [M+H]⁺.

EXAMPLE 7 5'-Fluoro-3'-(7-methyl-8-oxo-7,8-dihydroimidazo
[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

5'-Fluoro-3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile was coupled to give the title compound as an off-white solid (50 mg, 29%): δ_{H} (400 MHz, d₆-DMSO) 3.48 (3H, s), 7.22 (1H, d, J 6), 7.59-7.70 (5H, m), 7.79-7.87 (2H, m), 7.80 (1H, s), 8.03 (1H, dd, J 8 and 1); m/z (ES⁺) 345 [M+H]⁺.

EXAMPLE 8 2,2'-Difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo
30 [1,2-*a*]pyrazin-3-yl)-biphenyl-4-carbonitrile

2,2'-Difluoro-5'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-4-carbonitrile (prepared in the same way as in Example 19) was coupled to give the title compound as an off-white solid (30 mg, 16%): δ_{H} (400 MHz, d₆-DMSO) 3.48

(3H, s), 7.18 (1H, d, *J* 6), 7.59-7.61 (2H, m), 7.73 (1H, s), 7.75-7.82 (2H, m), 7.87-7.88 (2H, m), 8.05 (1H, d, *J* 10); *m/z* (ES⁺) 363 [M+H]⁺.

EXAMPLE 9 7-Ethyl-3-[4-fluoro-3-(2-methyl-2*H*-[1,2,4]triazol-3-ylmethoxy)phenyl]-7*H*-imidazo[1,2-*a*]pyrazin-8-one

3-Bromo-7-ethyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one and 4-fluoro-3-(2-methyl-2*H*-[1,2,4]triazol-3-ylmethoxy)benzeneboronic acid were coupled to give the title compound as a white solid (43 mg, 23%): δ_{H} (400 MHz, *d*₆-DMSO) 1.26 (3H, t, *J* 7), 3.94 (3H, s), 3.96 (2H, s), 5.52 (2H, s), 7.22-7.26 (2H, m), 7.44 (1H, dd, *J* 11 and 8), 7.55 (1H, d, *J* 6), 7.64 (1H, dd, *J* 8 and 2), 7.66 (1H, s), 7.98 (1H, s); *m/z* (ES⁺) 369 [M+H]⁺.

EXAMPLE 10 6,2'-Difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

6,2'-Difluoro-5'-(5,5-dimethyl-[1,3,2]dioxaborinan-2-yl)biphenyl-2-carbonitrile was coupled to produce the title compound: δ_{H} (400 MHz, CDCl₃) 3.59 (3H, s), 6.71 (1H, d, *J* 6.0), 7.37-7.67 (8H, m); *m/z* (ES⁺) 363 [M+H]⁺.

EXAMPLE 11 4,2'-Difluoro-5'-(7-ethyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

3-Bromo-7*H*-imidazo[1,2-*a*]pyrazin-8-one (prepared as described in Example 5, 1.5 g, 7 mmol) was suspended in *N,N*-dimethylformamide (15 ml) then treated with sodium hydride (420 mg of a 60% dispersion in mineral oil, 10.5 mmol). The resulting mixture was heated at 60°C for 20 min then treated with iodoethane (840 μ l, 10.5 mmol). After stirring for 40 min the reaction was cooled, diluted cautiously with water (150 ml) and then extracted with dichloromethane (2 x 100 ml). The organics were combined, washed with water, dried over anhydrous magnesium sulphate, filtered and concentrated under high vacuum. The residue was triturated with diethyl ether and the resulting solid collected by filtration and recrystallized from ethyl acetate/methanol to furnish 3-bromo-7-ethyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one as a white solid (0.34 g): δ_{H} (400 MHz, DMSO) 1.23 (3H, t, *J* 7), 3.94 (2H, q, *J* 7), 7.31 (1H, d, *J* 6), 7.41 (1H, d, *J* 6), 7.62 (1H, s); *m/z* (ES⁺) 244 and 242 [M+H]⁺.

Reaction of the above product under the same Suzuki coupling conditions as described in Example 5 gave the title compound after recrystallization from ethyl acetate/dichloromethane: δ_{H} (400 MHz, DMSO) 1.25 (3H, t, J 7.1), 3.96 (2H, q, J 7.1), 7.22 (1H, d, J 6.0), 7.57-7.61 (2H, m), 7.72 (1H, s), 7.76-7.84 (4H, m), 8.07 (1H, dd, J 8.6 and 2.7); m/z (ES^+) 377 $[\text{M}+\text{H}]^+$; Found C, 66.87; H, 3.54; N, 14.62. $\text{C}_{21}\text{H}_{14}\text{F}_2\text{N}_4\text{O}$ requires C, 67.02; H, 3.75; N, 14.89%.

Examples 12 to 18 were prepared in the same way as Example 10.

10 **EXAMPLE 12** 4,2'-Difluoro-5'-[7-(2,2-difluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

The title compound was prepared using 2-bromo-1,1-difluoroethane: δ_{H} (400 MHz, DMSO) 4.44 (2H, td, J 13 and 4.4), 6.36 (1H, tt, J 56 and 4.4), 7.19 (1H, d, J 6.1), 7.60-7.65 (2H, m), 7.74-7.86 (5H, m), 8.08 (1H, dd, J 8.6 and 2.7); m/z (ES^+) 413 $[\text{M}+\text{H}]^+$; Found C, 60.88; H, 2.97; N, 13.38. $\text{C}_{21}\text{H}_{12}\text{F}_4\text{N}_4\text{O}$ requires C, 61.17; H, 2.93; N, 13.59%.

EXAMPLE 13 4,2'-Difluoro-5'-[7-(1-methylethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

20 The title compound was prepared using 2-bromopropane: δ_{H} (400 MHz, DMSO) 1.33 (6H, d, J 6.8), 5.11 (1H, m), 7.20 (1H, d, J 6.1), 7.56-7.63 (2H, m), 7.74 (1H, m), 7.61-7.84 (4H, m), 8.06 (1H, dd, J 8.7 and 2.6); m/z (ES^+) 391 $[\text{M}+\text{H}]^+$; Found C, 68.00; H, 4.04; N, 14.03. $\text{C}_{22}\text{H}_{16}\text{F}_4\text{N}_4\text{O}$ requires C, 67.79; H, 4.13; N, 14.35%.

25 **EXAMPLE 14** 4,2'-Difluoro-5'-[7-(2-fluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

The title compound was prepared using 1-bromo-2-fluoroethane instead of iodoethane: δ_{H} (400 MHz, DMSO) 4.25 (1H, t, J 4.5), 4.31 (1H, t, J 4.5), 4.65 (1H, t, J 4.5), 4.77 (1H, t, J 4.5), 7.18 (1H, d, J 6.0), 7.59-7.62 (2H, m), 7.74 (1H, m), 7.75-7.85 (4H, m), 8.07 (1H, dd, J 8.7 and 2.6); m/z (ES^+) 395 $[\text{M}+\text{H}]^+$; Found C, 63.49; H, 3.27; N, 13.96. $\text{C}_{21}\text{H}_{13}\text{F}_3\text{N}_4\text{O}$ requires C, 63.96; H, 3.32; N, 14.21%.

EXAMPLE 15 4,2'-Difluoro-5'-[7-(2,2,2-trifluoroethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

The title compound was prepared using 1-iodo-2,2,2-trifluoroethane: δ_{H} (400 MHz, DMSO) 4.90 (2H, q, *J* 9.2), 7.20 (1H, d, *J* 6.0), 7.60-7.66 (2H, m), 7.75-7.85 (5H, m), 8.06 (1H, dd, *J* 8.7 and 2.6); *m/z* (ES⁺) 431 [M+H]⁺.

EXAMPLE 16 4,2'-Difluoro-5'-[7-(2-methyl-2*H*-[1,2,4]triazol-3-ylmethyl)-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

The title compound was prepared using 3-chloromethyl-2-methyl-2*H*-[1,2,4]triazole: δ_{H} (400 MHz, DMSO) 3.96 (3H, s), 5.35 (2H, s), 7.28 (1H, d, *J* 6.0), 7.62-7.64 (2H, m), 7.75-7.84 (6H, m), 8.06 (1H, dd, *J* 8.6 and 2.7); *m/z* (ES⁺) 444 [M+H]⁺; Found C, 62.48; H, 3.67; N, 21.98. C₂₃H₁₅F₂N₇O requires C, 62.30; H, 3.41; N, 22.11%.

EXAMPLE 17 4,2'-Difluoro-5'-[8-oxo-7-(pyridin-2-ylmethyl)-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl]biphenyl-2-carbonitrile

The title compound was prepared using 2-chloromethylpyridine hydrochloride: δ_{H} (400 MHz, DMSO) 5.26 (2H, s), 7.28-7.35 (3H, m), 7.61-7.63 (2H, m), 7.76-7.86 (6H, m), 8.06 (1H, dd, *J* 8.6 and 2.7), 8.50 (1H, m); *m/z* (ES⁺) 440 [M+H]⁺; Found C, 68.19; H, 3.48; N, 15.69. C₂₅H₁₅F₂N₅O requires C, 68.33; H, 3.44; N, 15.94%.

EXAMPLE 18 6,2'-Difluoro-5'-(7-ethyl-8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)-biphenyl-2-carbonitrile

This compound was prepared using 6,2'-difluoro-5'-(5,5-dimethyl-[1,3,2]dioxaborinan-2-yl)biphenyl-2-carbonitrile in the final step: δ_{H} (400 MHz, CDCl₃) 1.38 (3H, t, *J* 7.1), 4.04 (2H, q, *J* 7.1), 6.72 (1H, d, *J* 6.0), 7.38-7.67 (8H, m); *m/z* (ES⁺) 377 [M+H]⁺.

EXAMPLE 19 3-Chloro-2'-fluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-a]pyrazin-3-yl)biphenyl-4-carbonitrile

A mixture of 4-bromo-2-chlorobenzonitrile (2.2 g, 10 mmol) and 2-fluorobenzeneboronic acid (1.82 g, 13 mmol) in tetrahydrofuran (30 ml) was
5 treated with 2M sodium carbonate (13 ml) then degassed with nitrogen for 10 min. Tetrakis(triphenylphosphine)palladium(0) (0.58 g, 0.5 mmol) was added and this mixture was heated under reflux for 12 h. The resulting mixture was partitioned between dichloromethane and water. The organics were washed with water, brine, dried over anhydrous magnesium sulphate, filtered and
10 concentrated *in vacuo*. Purification of the residue by chromatography on silica gel eluting with isohexane on a gradient of ethyl acetate (5-15%) gave 3-chloro-2'-fluorobiphenyl-4-carbonitrile as a white solid (2.3 g, 99%): δ_H (400 MHz, $CDCl_3$) 7.17-7.57 (5H, m), 7.64-7.75 (2H, m).

To a slurry of the above product (2.3 g, 10 mmol) and 1,3-dibromo-5,5-dimethylhydantoin (3.1 g, 11 mmol) in acetonitrile (20 ml) was added
15 concentrated sulphuric acid (0.8 ml). The slurry was warmed to 70°C and the resulting solution stirred at 70°C for 16 h. After 16 h, the reaction was complete. Water (75 ml) was added and the resulting suspension was filtered and dried, to give 5'-bromo-3-chloro-2'-fluorobiphenyl-4-carbonitrile as an off-white solid (3.1 g,
20 98%): δ_H (360 MHz, $CDCl_3$) 7.07-7.17 (1H, m), 7.29-7.76 (5H, m).

The above product (0.6 g, 2 mmol), 4,4,5,5,4',4',5',5'-octamethyl-[2,2']bi[[1,3,2]dioxaborolanyl] (0.6 g, 2.4 mmol) and potassium acetate (0.39 g, 4 mmol) were suspended in 1,4-dioxane (6 ml) and the mixture was degassed with nitrogen for 10 minutes, before adding 1,1'-bis(diphenylphosphino)ferrocene-dichloropalladium(II) complex with dichloromethane (0.05 g, 0.06 mmol). The
25 resulting mixture was heated at 90°C for 12 h. The mixture was cooled to ambient temperature, filtered and the filter-cake washed with dichloromethane. The filtrate was evaporated to dryness and the residue stirred with ice-cold 2N sodium hydroxide (70 ml) for 20 minutes. The aqueous mixture was filtered and
30 the filtrate washed with diethyl ether (2 x 50 ml). The organics were discarded and the aqueous phase cooled to 0°C before lowering the pH to 6 by addition of 36% hydrochloric acid. The product was extracted into diethyl ether to give 3-chloro-2'-fluoro-5'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-4-carbonitrile as a yellow oil (0.53 g, 74%).

3-Bromo-7-methyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one and the above product were coupled in the same way as in Example 5 to give the title compound as an off-white solid (50 mg, 26%): δ_H (360 MHz, d_6 -DMSO) 3.48 (3H, s), 7.18 (1H, d, *J* 6), 7.56-7.59 (2H, m), 7.70-7.90 (2H, m), 8.12 (2H, dd, *J* 8 and 3); *m/z* (ES⁺) 379 [M+H]⁺.

EXAMPLE 20 3-[4-Fluoro-3-(1-methyl-1*H*-[1,2,3]triazol-4-ylmethoxy)phenyl]-7-methyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one

A cooled (-20°C) solution of 2,2,6,6-tetramethylpiperidine (28 ml, 165 mmol) in tetrahydrofuran (400 ml) was treated with *n*-butyllithium (63 ml of a 2.5M solution in hexanes, 157.5 mmol). This mixture was then cooled to -78°C. 1-Bromo-4-fluorobenzene (16.5 ml, 150 mmol) was then added neat and dropwise over 10 min and stirring at -78°C was continued for 3 h. Triisopropyl borate (40 ml, 172.5 mmol) was then added and stirring at -78°C continued for 30 min before removing the cooling bath. When the internal temperature of the reaction reached -40°C, 5N hydrochloric acid was added (75 ml) and the mixture was stirred to ambient temperature. After stirring at ambient temp for 1 h the majority of the tetrahydrofuran was removed and the mixture partitioned between ether (500 ml) and 1N hydrochloric acid (500 ml). The organics were then extracted with 2N sodium hydroxide (400 ml) and the organics were discarded. The aqueous was cooled in an ice-water bath and 5N hydrochloric acid (150 ml) was added dropwise over 15 min. The resulting white solid was collected and dried under vacuum to afford 5-bromo-2-fluorobenzeneboronic acid (25 g, 76%).

A solution of the above product (25 g, 114 mmol) in tetrahydrofuran was treated with hydrogen peroxide (7.8 ml of a 35 wt % solution in water) then with sodium hydroxide (1.4 ml of a 4N solution in water). A mild exotherm caused the internal temperature to reach 40°C. This mixture was left to stir at ambient temperature for 14 h then treated with manganese dioxide (200 mg) and stirring was continued for 90 min before filtering the reaction (GF/A filter paper). The filtrate was concentrated on a rotary evaporator and the residue partitioned between ether (400 ml) and water. The organics were washed with more water, brine and dried over anhydrous magnesium sulphate. Filtration and evaporation

to dryness afforded 5-bromo-2-fluorophenol (19.7 g, 90%) as a colourless liquid: δ_H (400 MHz, d_6 -DMSO) 6.93-6.97 (1H, m), 7.09-7.14 (2H, m), 10.36 (1H, br).

An ice-cooled solution of the above product (1.91 g, 10 mmol), (1-methyl-1*H*[1,2,3]triazol-4-yl)methanol (1.24 g, 11 mmol) and triphenylphosphine (3.93 g, 15 mmol) in tetrahydrofuran (50 ml) was treated dropwise with diisopropyl azodicarboxylate (3.03 g, 15 mmol) over 10 min. The resulting mixture was stirred to ambient temperature over 12 h and then glacial acetic acid (1 ml) was added. The reaction mixture was concentrated *in vacuo* then partitioned between ethyl acetate and 0.01N sodium hydroxide solution. The organics were washed with water, brine, dried over anhydrous magnesium sulphate and concentrated to give an oil. This oil was purified by chromatography on silica gel eluting with isohexane on a gradient of ethyl acetate (20-50%). Product-containing fractions were concentrated and the resulting residue triturated with 10% ether in isohexane to furnish 4-(5-bromo-2-fluorophenoxymethyl)-1-methyl-1*H*[1,2,3]triazole as a white solid (1.9 g, 66%): δ_H (360 MHz, d_6 -DMSO) 4.06 (3H, s), 5.25 (2H, s), 7.10-7.20 (2H, m), 7.55-7.70 (1H, m), 8.19 (1H, s).

The above product was converted to 4-fluoro-3-(1-methyl-1*H*[1,2,3]triazol-4-ylmethoxy)benzeneboronic acid as described in Example 19: δ_H (400 MHz, d_6 -DMSO) 4.06 (3H, s), 5.19 (2H, s), 7.16 (1H, dd, *J* 12 and 8), 7.38-7.41 (1H, m), 7.71 (1H, dd, *J* 9 and 1), 8.09 (2H, s), 8.16 (1H, s).

The above product was coupled as described in Example 5 to afford the title compound as a white solid: δ_H (400 MHz, d_6 -DMSO) 3.48 (3H, s), 4.06 (3H, s), 5.35 (2H, s), 7.17-7.21 (2H, m), 7.39 (1H, dd, *J* 11 and 8), 7.55 (1H, d, *J* 6), 7.60 (1H, dd, *J* 8 and 2), 7.67 (1H, s), 8.22 (1H, s); *m/z* (ES^+) 355 [$M+H$] $^+$.

EXAMPLE 21 3-[4-Fluoro-3-(2-methyl-2*H*[1,2,4]triazol-3-ylmethoxy)phenyl]-7-methyl-7*H*imidazo[1,2-*a*]pyrazin-8-one

A cooled (0°C) solution of 5-bromo-2-fluorophenol, (2-methyl-2*H*[1,2,4]triazol-3-yl)methanol and triphenylphosphine in tetrahydrofuran was treated dropwise with diisopropyl azodicarboxylate over 10 min. The resulting mixture was stirred overnight and then acetic acid was added. The reaction mixture was evaporated to dryness and the residue was partitioned between ethyl acetate and 0.01N sodium hydroxide solution. The organic phase was washed with water and brine, dried over anhydrous magnesium sulphate and

pre-adsorbed onto silica. Purification by flash column chromatography on silica gel, eluting with isohexane on a gradient of ethyl acetate (10-50%), led to product-containing fractions that were combined. The material was further purified by flash column chromatography on silica gel, eluting with dichloromethane containing 1% ammonia on a gradient of methanol (0.5-3%), to give 5-(5-bromo-2-fluoro-phenoxy-methyl)-1-methyl-1*H*-[1,2,4]triazole as an off-white solid (2.1 g, 37%): δ_H (360 MHz, $CDCl_3$) 4.01 (3H, s), 5.29 (2H, s), 6.96-6.99 (1H, m), 7.09-7.13 (1H, m), 7.29 (1H, dd, *J* 5 and 2), 7.88 (1H, s); *m/z* (ES^+) 288 [MH^+].

The above product was reacted with 4,4,5,5,4',4',5',5'-octamethyl-[2,2']bi[[1,3,2]-dioxaborolanyl] as described in Example 19 to give the title compound as a pale yellow solid (0.43 g, 62%).

The above product was coupled in the same way as in Example 5 to give the title compound as a white solid (58 mg, 33%): δ_H (400 MHz, d_6 -DMSO) 3.48 (3H, s), 3.94 (3H, s), 5.52 (2H, s), 7.21-7.25 (2H, m), 7.45 (1H, dd, *J* 11 and 8), 7.53 (1H, d, *J* 6), 7.62 (1H, dd, *J* 8 and 2), 7.65 (1H, s), 7.98 (1H, s); *m/z* (ES^+) 355 [$M+H$] $^+$.

EXAMPLE 22 4,2'-Difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

A suspension of 8-chloroimidazo[1,2-*a*]pyrazine (5 g, 325 mmol) in 5*N* hydrochloric acid (35 ml) was heated at 90°C for 14 h. The reaction mixture was cooled, then concentrated using rotary evaporation. Azeotropic removal of water with toluene (2 x 30 ml) gave 8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazine hydrochloride (5.7 g): δ_H (360 MHz, DMSO) 7.01 (1H, d, *J* 5.6), 7.60 (1H, d, *J* 5.6), 7.70 (1H, s), 7.97 (1H, s), 11.59 (1H, br s).

The above product (1 g) was suspended in *N,N*-dimethylformamide (15 ml) then treated with sodium hydride (650 mg of a 60% dispersion in mineral oil). The resulting mixture was heated at 60°C for 20 min then cooled to room temperature and treated with iodomethane (0.4 ml). After stirring for 15 h the reaction mixture was diluted cautiously with methanol (5 ml) and then silica gel (5 g) was added and the solvents removed under vacuum. The silica residue was filtered through a plug of silica gel, using 10% methanol in dichloromethane as eluent, then the solvents were removed *in vacuo* to leave 7-methyl-7*H*-imidazo[1,2-*a*]pyrazin-8-one (1.15 g): δ_H (360 MHz, DMSO) 3.43 (3H, s), 7.11 (1H,

d, J 5.8), 7.46 (1H, s), 7.56 (1H, d, J 5.8), 7.78 (1H, s). NOE studies showed that irradiation of the 3H singlet at δ 3.43 gave enhancement of the doublet at δ 7.11 and vice-versa indicating that methylation took place on the 7-nitrogen rather than the 8-oxygen.

5 The above product (315 mg, 2.1 mmol) in a Parr flask was dissolved in methanol (20 ml) and 10% Pd-C catalyst (100 mg) was added. The reaction mixture was shaken under 55 psi of hydrogen for 16 days then filtered and concentrated under vacuum to leave a residue. This was recrystallized from ethyl acetate/methanol to give 7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazine
10 (185 mg): δ_H (360 MHz, DMSO) 3.31 (3H, s), 3.70-3.73 (2H, m), 4.24-4.28 (2H, m), 7.08 (1H, s), 7.32 (1H, s).

 The above product (80 mg, 0.53 mmol) and 5'-bromo-4,2'-difluorobiphenyl-2-carbonitrile (prepared as described in WO 02/074773, 204 mg, 1.3 mol eq) were dissolved in DMA (2 ml) and potassium acetate (280 mg, 1.5 mol eq), palladium
15 acetate (6 mg, 5 mol %) and triphenylphosphine (7 mg, 5 mol %) were all added. After bubbling nitrogen through the reaction mixture for 10 min, the temperature was raised to 130°C and heating continued for 3 h. After cooling, the crude reaction mixture was poured onto a column of silica and purified using 0-5% methanol in dichloromethane as eluent. Recrystallization from ethyl
20 acetate/dichloromethane gave the title compound (18 mg), mp 224°C: δ_H (400 MHz, DMSO) 3.02 (3H, s), 3.72-3.75 (2H, m), 4.35-4.38 (2H, m), 7.38 (1H, s), 7.57 (1H, m), 7.72-7.78 (4H, m), 8.06 (1H, dd, J 9 and 3); m/z (ES⁺) 365 [M+H]⁺.

 Examples 23 and 24 were prepared by the method of Example 22:

25

EXAMPLE 23 5,2'-Difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo
[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

 The title compound was prepared using 5'-bromo-5,2'-difluorobiphenyl-2-carbonitrile in the final step: δ_H (400 MHz, CDCl₃) 3.18 (3H, s), 3.75-3.77 (2H, m),
30 4.32-4.34 (2H, m), 7.24-7.28 (1H, m), 7.30-7.38 (3H, m), 7.44-7.46 (1H, m), 7.52-7.55 (1H, m), 7.83 (1H, dd, J 7 and 4); m/z (ES⁺) 365 [M+H]⁺.

EXAMPLE 24 6,2'-Difluoro-5'-(7-methyl-8-oxo-5,6,7,8-tetrahydroimidazo
[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

The title compound was prepared using 5'-bromo-6,2'-difluorobiphenyl-2-carbonitrile in the final step: δ_H (400 MHz, $CDCl_3$) 3.19 (3H, s), 3.73-3.78 (2H, m),
5 4.28-4.35 (2H, m), 7.34-7.42 (3H, m), 7.45-7.49 (1H, m), 7.54-7.58 (2H, m), 7.64
(1H, m); m/z (ES^+) 365 $[M+H]^+$.

EXAMPLE 25 4,2'-Difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo
[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

10 8-Oxo-7,8-dihydroimidazo[1,2-*a*]pyrazine hydrochloride (see Example 20, 1
g) in a Parr flask was dissolved in methanol (60 ml) and 10% Pd-C catalyst (300
mg) was added. The reaction mixture was shaken under 55 psi of hydrogen for 17
days then filtered and concentrated under vacuum to leave a residue. This was
recrystallized from ethyl acetate/methanol to give 8-oxo-5,6,7,8-
15 tetrahydroimidazo[1,2-*a*]pyrazine hydrochloride (610 mg): δ_H (360 MHz, DMSO)
3.66-3.70 (2H, m), 4.36-4.39 (2H, m), 7.67 (1H, s), 7.78 (1H, s), 8.84 (1H, br s).

The above product (500 mg, 40 mmol) was suspended in *N,N*-
dimethylformamide (10 ml) then treated with sodium hydride (380 mg of a 60%
dispersion in mineral oil, 2.4 mol eq). The resulting mixture was heated at 60°C
20 for 20 min then cooled to room temperature and treated with iodoethane (0.386
ml, 1.2 mol eq). After stirring for 15 h the reaction mixture was diluted
cautiously with methanol (5 ml) and then silica gel (4 g) was added and the
solvents removed under vacuum. The silica residue was filtered through a plug
of silica gel, using 10% methanol in dichloromethane as eluent, then the solvents
25 were removed *in vacuo* to leave 7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-
a]pyrazine (400 mg): δ_H (400 MHz, DMSO) 1.10 (3H, t, J 7.2), 3.47 (2H, J 7.2),
3.70-3.73 (2H, m), 4.23-4.27 (2H, m), 7.08 (1H, s), 7.32 (1H, s).

The above product (82.5 mg, 0.53 mmol) and 5'-bromo-4,2'-
difluorobiphenyl-2-carbonitrile (prepared as described in WO 02/074773, 204 mg,
30 1.3 mol eq) were dissolved in DMA (2 ml) and potassium acetate (280 mg, 1.5 mol
eq), palladium acetate (6 mg, 5 mol %) and triphenylphosphine (7 mg, 5 mol %)
were all added. After bubbling nitrogen through the reaction mixture for 10 min,
the temperature was raised to 130°C and heating continued for 3 h. After
cooling, the crude reaction mixture was poured onto a column of silica and

purified using 0-5% methanol in dichloromethane as eluent. Recrystallization from ethyl acetate/dichloromethane gave the title compound (18 mg): δ_{H} (400 MHz, DMSO) 1.12 (3H, t, J 7.1), 3.51 (2H, q, J 7.1), 3.73-3.76 (2H, m), 4.34-4.37 (2H, m), 7.39 (1H, s), 7.57 (1H, m), 7.72-7.78 (4H, m), 8.06 (1H, dd, J 9 and 3); m/z (ES⁺) 379 [M+H]⁺.

Examples 26 to 28 were prepared using the method of Example 25 using 5'-bromo-5,2'-difluorobiphenyl-2-carbonitrile and 5'-bromo-6,2'-difluorobiphenyl-2-carbonitrile respectively (prepared as described in WO 02/074773):

10

EXAMPLE 26 5,2'-Difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

This compound was prepared using 5'-bromo-5,2'-difluorobiphenyl-2-carbonitrile (prepared as described in WO 02/074773): δ_{H} (400 MHz, DMSO) 1.11 (3H, t, J 7.1), 3.51 (2H, q, J 7.1), 3.73-3.76 (2H, m), 4.35-4.38 (2H, m), 7.40 (1H, s), 7.55-7.61 (2H, m), 7.68-7.78 (3H, m), 8.13 (1H, dd, J 9 and 3); m/z (ES⁺) 379 [M+H]⁺.

20

EXAMPLE 27 6,2'-Difluoro-5'-(7-ethyl-8-oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

δ_{H} (400 MHz, DMSO) 1.20 (3H, t, J 7.1), 3.59 (2H, q, J 7.1), 3.82-3.85 (2H, m), 4.39-4.45 (2H, m), 7.45 (1H, s), 7.65-7.70 (1H, m), 7.83-7.92 (4H, m), 8.00 (1H, dd, J 9 and 3); m/z (ES⁺) 379 [M+H]⁺.

25

EXAMPLE 28 4,2'-Difluoro-5'-[8-oxo-7-(pyridin-2-ylmethyl)-5,6,7,8-tetrahydro-imidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile

30

8-Oxo-5,6,7,8-tetrahydroimidazo[1,2-*a*]pyrazine hydrochloride (87 mg, 0.53 mmol), prepared as described in Example 25, was reacted with 5'-bromo-4,2'-difluorobiphenyl-2-carbonitrile (prepared as described in WO 02/074773, 204 mg, 1.3 mol eq) in DMA (2 ml) in the presence of potassium acetate (280 mg, 1.5 mol eq), palladium acetate (6 mg, 5 mol %) and triphenylphosphine (7 mg, 5 mol %). After bubbling nitrogen through the reaction mixture for 10 min, the temperature was raised to 130°C and heating continued for 3 h. After cooling, the crude

reaction mixture was poured onto a column of silica and purified using 0-5% methanol in dichloromethane as eluent. Recrystallization from ethyl acetate/dichloromethane gave 4,2'-difluoro-5'-(8-oxo-7,8-dihydroimidazo[1,2-*a*]pyrazin-3-yl)biphenyl-2-carbonitrile (126 mg): δ_{H} (400 MHz, DMSO) 3.54-3.58 (2H, m), 4.28-4.32 (2H, m), 7.39 (1H, s), 7.54-7.59 (1H, m), 7.71-7.78 (4H, m), 8.05-8.08 (1H, m), 8.18 (1H, br s); m/z (ES^+) 351 $[\text{M}+\text{H}]^+$.

The title compound was prepared using the method described in the first part of Example 11, using 2-chloromethylpyridine hydrochloride: δ_{H} (400 MHz, DMSO) 3.60-3.64 (2H, m), 4.18-4.22 (2H, m), 4.60 (2H, m), 7.08-7.11 (1H, m), 7.16-7.19 (1H, m), 7.21 (1H, s), 7.54-7.59 (1H, m), 7.52-7.56 (5H, m), 7.82-7.86 (1H, m), 8.30-8.32 (1H, m); m/z (ES^+) 442 $[\text{M}+\text{H}]^+$.

EXAMPLE 29 4,2'-Difluoro-5'-(7-methyl-8-oxo-7,8-dihydroimidazo[1,2-*d*][1,2,4]triazin-3-yl)biphenyl-2-carbonitrile

2-Hydrazinocarbonylimidazole (prepared according to the procedure described in EP-A-0713876) (1.00 g, 7.93 mmol) was suspended in Dowtherm A (6 ml), *N,N*-dimethylformamide dimethyl acetal (1.89 g, 15.9 mmol) added and the mixture heated at 100°C for 1.5 h then at 210°C for 18 h. The mixture was allowed to cool to room temperature, diethyl ether (100 ml) added and the precipitated solid was filtered. The brown solid was purified by dry flash column chromatography on silica eluting with dichloromethane (+0.1% 0.880 ammonia) on a gradient of methanol (5-12%). Collecting appropriate fractions gave 7*H*-imidazo[1,2-*d*][1,2,4]triazin-8-one (216 mg, 20%) as an off-white solid: δ_{H} (400 MHz, CDCl_3) 7.59 (1H, d, *J* 1), 7.91 (1H, d, *J* 1), 8.74 (1H, s), 12.39 (1H, s).

Sodium hydride (60% suspension in mineral oil) (76 mg, 1.90 mmol) was added to a suspension of the above product (216 mg, 1.59 mmol) in *N,N*-dimethylformamide (4 ml) and the mixture stirred at 20°C for 10 min then at 60°C for 5 min. The mixture was allowed to cool to ambient temperature, methyl iodide (676 mg, 4.76 mmol) added and the mixture left to stir for 10 min. The solvent was evaporated and the residue was purified by flash column chromatography on silica eluting with dichloromethane on a gradient of methanol (2-4%). Collecting appropriate fractions gave 7-methyl-7*H*-imidazo[1,2-*d*][1,2,4]triazin-8-one (150 mg, 63%) as a pale yellow solid: δ_{H} (360 MHz, CDCl_3) 7.45 (1H, d, *J* 1), 7.63 (1H, d, *J* 1), 8.19 (1H, s).

The above product (75 mg, 0.50 mmol), 5'-bromo-4,2'-difluorobiphenyl-2-carbonitrile (prepared according to the procedure described in WO 02/38568) (220 mg, 0.75 mmol) and potassium acetate (74 mg, 0.75 mmol) were suspended in *N,N*-dimethylacetamide (2 ml) and degassed with nitrogen for 10 min.

5 Palladium(II) acetate (5.6 mg, 0.025 mmol) and triphenylphosphine (6.6 mg, 0.025 mmol) were added and the mixture heated at 130°C for 2 h. The mixture was allowed to cool to ambient temperature, diluted with ethyl acetate (100 ml) and washed with water (50 ml) and brine (50 ml), dried over anhydrous sodium sulfate, filtered and evaporated to give a yellow solid. The solid was purified by
10 flash column chromatography on silica eluting with 2% methanol in dichloromethane. Collecting appropriate fractions followed by trituration with diethyl ether gave the title compound as a white solid: δ_H (400 MHz, $CDCl_3$) 3.80 (3H, s), 7.41-7.47 (2H, m), 7.53-7.64 (4H, m), 7.67 (1H, s), 8.42 (1H, s); m/z (ES⁺) 364 (M⁺+H).

15

EXAMPLE 30 3'-(1-Methyl-6-oxo-1,6-dihydropurin-9-yl)biphenyl-2-carbonitrile

Sodium methoxide (2.70 g, 50.0 mmol) was added portionwise over 10 min to a 0°C suspension of 4,6-dichloro-5-nitropyrimidine (4.85 g, 25.0 mmol) in
20 methanol (90 ml). On complete addition the mixture was stirred at 0°C for 2 h. The precipitate was filtered and the filtrate evaporated. The residue was suspended in isohexane and filtered. The filtrate was evaporated and the residue purified by flash column chromatography on silica, eluting with 5% ethyl acetate in isohexane. Collecting appropriate fractions gave 4-chloro-6-methoxy-5-
25 nitropyrimidine (3.50 g, 74%) as a white solid: δ_H (400 MHz, $CDCl_3$) 4.15 (3H, s), 8.65 (1H, s).


The above product (3.50 g, 18.5 mmol) and 3'-aminobiphenyl-2-carbonitrile (prepared according to the procedure described in WO 02/38568) (3.23 g, 16.6 mmol) were dissolved in triethylamine (9.32 g, 92.3 mmol) and dimethylsulfoxide
30 (10 ml) and heated at 90°C for 18 h. The mixture was allowed to cool to ambient temperature then poured onto water (150 ml) and extracted with ethyl acetate (2 x 300 ml). The organic fractions were combined, washed with water (3 x 75 ml) and brine (50 ml), dried over anhydrous sodium sulfate, filtered and evaporated to give a brown solid. The brown solid was triturated with diethyl ether (50 ml),

filtered, washed with diethyl ether (2 x 10 ml) and left to air dry, which gave 3'-(1-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-4-ylamino)biphenyl-2-carbonitrile (2.24 g, 38%) as a yellow solid: δ_H (360 MHz, $CDCl_3$) 3.52 (3H, s), 7.45-7.57 (5H, m), 7.68 (1H, ddd, J 8, 8 and 1), 7.78-7.82 (2H, m), 8.04 (1H, s), 11.13 (1H, s).

- 5 The above product (100 mg, 0.28 mmol) was suspended in THF (5 ml) and ethanol (5 ml), tin(II) chloride (162 mg, 0.72 mmol) added and the mixture heated under reflux for 18 h. The solvent was evaporated and the residue suspended in formic acid (20 ml) and heated under reflux for 48 h. The mixture was allowed to cool to ambient temperature and the solvent evaporated. The residue was
- 10 dissolved in dichloromethane and methanol and evaporated onto silica. The product was purified by flash column chromatography on the same, eluting with dichloromethane (+0.1% 0.880 ammonia) on a gradient of methanol (2-5%). Collecting appropriate fractions gave the title compound (35 mg, 37%) as a white
- 15 solid: δ_H (500 MHz, DMSO) 3.55 (3H, s), 7.65 (1H, dd, J 8 and 8), 7.72-7.79 (3H, m), 7.85 (1H, ddd, J 8, 8 and 1), 7.97 (1H, dd, J 8 and 1), 8.00-8.04 (2H, m), 8.45 (1H, s), 8.59 (1H, s); m/z (ES⁺) 328 (M⁺+H).

CLAIMS

1. A compound of formula I, or a pharmaceutically acceptable salt thereof:



(I)

wherein

-U-V- represents -CH=CH-, or -CH₂-CH₂-, -N=CH- or -CH=N-;

X¹ represents hydrogen, halogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy;

10 X² represents hydrogen or halogen;

Y represents a chemical bond, an oxygen atom, or a -NH- or -OCH₂- linkage;

Z represents an optionally substituted aryl or heteroaryl group;

15 R¹ represents hydrocarbon, a heterocyclic group, trifluoromethyl, -SO₂R^a, -SO₂NR^aR^b, -COR^a, -CO₂R^a or -CONR^aR^b; and

R^a and R^b independently represent hydrogen, hydrocarbon or a heterocyclic group.

2. A compound according to claim 1 in which -U-V- represents -CH=CH- and
20 Y represents a chemical bond, an oxygen atom, or an -NH- linkage.

3. A compound according to claim 1 or 2 in which Y is a chemical bond.

4. A compound according to claim 1, 2 or 3 in which R¹ is methyl, ethyl,
25 fluoroethyl, difluoroethyl, trifluoroethyl, isopropyl, methyl-triazolylmethyl or pyridinylmethyl.

5. A compound according to any preceding claim in which X¹ is fluorine and X² is hydrogen.
6. A compound according to any preceding claim in which Z is a substituted phenyl or triazolyl group.
7. A pharmaceutical composition comprising a compound according to any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
8. A compound according to any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof, for use in a method of treatment of the human body by therapy.
9. Use of a compound according to any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for treating or preventing anxiety, convulsions or a cognitive disorder.
10. A method for the treatment and/or prevention of anxiety, convulsions or a cognitive disorder which comprises administering to a patient in need of such treatment an effective amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof.

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 03/04685

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/4985 A61K31/53 A61P25/00
/(C07D487/04,241:00,235:00),(C07D487/04,253:00,235:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02 10170 A (MERCK SHARP & DOHME (GB)) 7 February 2002 (2002-02-07) cited in the application claims 1,8,9	1,9
A	WO 02 38568 A (MERCK SHARP & DOHME (GB)) 16 May 2002 (2002-05-16) claims 1,15	1,9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

29 January 2004

Date of mailing of the international search report

05/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

Internat'l application No.
PCT/GB 03/04685

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 10 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 03/04685

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0210170	A	07-02-2002	AU 7085301 A	13-02-2002
			CA 2417121 A1	07-02-2002
			EP 1307456 A1	07-05-2003
			WO 0210170 A1	07-02-2002
WO 0238568	A	16-05-2002	AU 1253002 A	21-05-2002
			CA 2427779 A1	16-05-2002
			EP 1343788 A1	17-09-2003
			WO 0238568 A1	16-05-2002
			WO 03008418 A1	30-01-2003
			US 2003055060 A1	20-03-2003
			US 2003216395 A1	20-11-2003

THIS PAGE BLANK (USPTO)